

Laboratory Studies of Carbon Kinetic Isotope Effects on the Production Mechanism of Particulate Phenolic Compounds Formed by Toluene Photooxidation: A Tool To Constrain Reaction Pathways


Satoshi Irei,^{*,†,§} Jochen Rudolph,[†] Lin Huang,[‡] Janeen Auld,^{†,||} Fabrice Collin,^{†,⊥} and Donald Hastie[†]

[†]Centre for Atmospheric Chemistry and Department of Chemistry, York University, 4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada

[‡]Climate Research Division, Atmospheric Science and Technology Directorate, Science and Technology Branch, Environment Canada, 4905 Dufferin Street, Toronto, Ontario, M3H 5T4, Canada

S Supporting Information

ABSTRACT: In this study, we examined compound-specific stable carbon isotope ratios for phenolic compounds in secondary organic aerosol (SOA) formed by photooxidation of isotope-label-free toluene. SOA generated by photooxidation of toluene using a continuous-flow reactor and an 8 m³ indoor smog chamber was collected on filters, which were extracted with acetonitrile for compound-specific analysis. Eight phenolic compounds were identified in the extracts using a gas chromatograph coupled with a mass spectrometer, and their compound-specific stable carbon isotope ratios were determined using a gas chromatograph coupled with a combustion furnace followed by an isotope ratio mass spectrometer. The majority of products, including methylnitrophenols and methylnitrocatechols, were isotopically depleted by 5–6‰ compared to the initial isotope ratio of toluene, whereas the isotope ratio for 4-nitrophenol remained identical to that of toluene. On the basis of the reaction mechanisms proposed in previous reports, stable carbon isotope ratios of these products were calculated. By comparing the observed isotope ratios with the predicted isotope ratios, we explored possible production pathways for the



particulate phenolic compounds.

1. INTRODUCTION

Airborne particulate organic matter or organic aerosol (OA), which is a predominant component of atmospheric aerosols,¹ has been of great interest in the past two decades owing to its possible association with adverse health effects² and with the global budget of radiative forcing.³ However, these associations are not clearly understood. One issue is the limited quantitative information available on secondary OA (SOA), which is formed by oxidative conversion of volatile organic compounds in the atmosphere. These difficulties are attributed to an insufficient understanding of chemical reaction mechanisms leading to the formation of SOA. A newly developed approach that can provide novel insights into SOA formation is studying the

ment provides useful indications regarding atmospheric photochemistry.^{5–9} One advantage of this technique is that the measurements do not require labeled reactants with stable isotopes. Instead, relatively small changes in natural isotope abundance are used to evaluate the extent of reaction processing. This allows isotope effects determined from laboratory studies to be directly applied to ambient observations. A kinetic isotope effect (KIE), which is an isotope fractionation during a chemical reaction, is reaction dependent; thus, measuring changes in isotope ratios during chemical reactions will provide additional insights into reaction mechanisms, such as SOA formation. To our knowledge, few studies have examined isotope fractionation during SOA

developed compound-specific stable carbon isotope analysis for nitrophenols.¹³

Nitro and nitrohydroxy aromatic compounds have been detected in gas-phase products^{14,15} as well as SOA^{16–19} produced by reactions of aromatic hydrocarbons with OH radicals in laboratory studies. On the basis of these laboratory studies, production mechanisms have been postulated. Thus, the application of compound-specific label-free isotope measurement to these compounds can be used to constrain reaction channels in reaction mechanism studies. In this report, the results are discussed based on postulated processes for the formation of phenolic products with the estimated carbon KIEs for the individual reaction steps.

2. EXPERIMENTAL SECTION

2.1. Flow Reactor and Smog Chamber Experiments.

Laboratory studies of SOA formed by the photooxidation of toluene were conducted with a continuous-flow reactor and a smog chamber. The former has features that production of SOA is theoretically unlimited, but the extent of toluene reaction is limited due to the low residence time. Meanwhile, the latter apparatus has features that the extent of toluene reaction goes on as long as UV lights are on (i.e., capability of high toluene turnover), but production of SOA is limited due to the limited volume of the apparatus. Details of the flow reactor and smog chamber experiments were described previously.^{10,11,19,20} Briefly, a mixture of synthetic air, NO (BOC gases), toluene (purity >99%, Sigma-Aldrich Canada), and synthesized isopropyl nitrite was irradiated with UV light to generate OH radicals. The initial mixing ratios of NO_x, toluene, and isopropyl nitrite (IPN) were approximately 5, 20–40, and 230 ppmv for the flow reactor experiments and 3.5, 0.5, and 0.7 ppmv for the smog chamber experiments, respectively. The initial $\delta^{13}\text{C}$ values of toluene determined based on conventional off-line isotope ratio measurements were $-26.72\text{‰} \pm 0.01\text{‰}$ and $-27.05\text{‰} \pm 0.02\text{‰}$ for the flow reactor experiments and the smog chamber experiments, respectively. Although toluene used for these experiments was from the same batch, the slightly heavier $\delta^{13}\text{C}$ for the flow reactor experiments (i.e., -26.72‰) was due to the small isotope fractionation during partial evaporation from bulk toluene. A detailed description of the evaporative isotope fractionation was discussed previously.^{10,21} This isotope fractionation is only 1/20 of the isotope fractionation observed for the toluene reaction with OH radicals, and did not significantly impact our observations. No seed particles were introduced in the flow reactor experiments, while dried ammonium sulfate particles were introduced as SOA seeds in the smog chamber experiments (3×10^4 particles cm^{-3} with a monomodal distribution and a maximum diameter

chromatograph coupled with a flame ionization detector. In the flow reactor and smog chamber experiments, the SOA was collected on 25 mm o.d. and 47 mm o.d. polytetrafluoroethylene (PTFE)-coated glass fiber disk filters (Pallflex Fiberfilm; Pall Corp., Port Washington, NY), respectively. The filters were used without precleaning. Filter sampling is summarized in Table 1. Sampling flow rates for the flow reactor

Table 1. Summary of Initial Experimental Conditions and SOA Filter Sampling

	FR1–FR5 ^a	SCI–SCIII ^b
initial experimental conditions		
NO _x (ppmv)	5	3.5
toluene (ppmv)	20–40	0.5
IPN (ppmv)	230	0.7
seed particles (particles cm^{-3})	0	30000
filter sampling		
number of filter samples collected	5	3
sampling duration (h)	2.4–19.0	5.03–5.22
sampling flow rate (L min^{-1})	6.3	12
sampled volume (m^{-3})	0.1–0.9	3.6–3.8

^aFlow reactor experiments. ^bSmog chamber experiments.

and the smog chamber experiments were 6.3 and 12 L min^{-1} , respectively. These corresponded to linear velocities of 1700 and 1300 cm min^{-1} , respectively. These linear velocities were smaller than that used for usual high-volume sampling (~ 2000 cm min^{-1}); therefore, the blow-off effect of collected SOA was lower than that for typical ambient particulate filter sampling using a high-volume air sampler.

2.2. Analysis of Filter Samples. Briefly, 5 or 24 μg carbon ($\mu\text{g C}$) of catechol, 9 or 26 $\mu\text{g C}$ of 2-nitrophenol, 6 or 23 $\mu\text{g C}$ of 4-ethylresorcinol, and 52 or 16 $\mu\text{g C}$ of 2,6-dimethyl-4-nitrophenol (>99% purity; Sigma-Aldrich, Oakville, ON, Canada) were spiked to a filter sample as internal standards (IS) for quantitative analysis of phenolic compounds and routine isotope measurements. Hereafter, these internal standards are referred to as IS1, IS2, IS3, and IS4, respectively. The filter was then extracted with 2–3 mL of HPLC-grade acetonitrile (Sigma-Aldrich, Oakville, ON, Canada) three times by sonication for 3–5 min in a 60 mL glass jar (wide-mouth amber glass jars; Chromatographic Specialties Inc., Brockville, ON, Canada) presilanized with dimethyldichlorosilane (Sigma-Aldrich, Oakville, ON, Canada). The extracts were filtered using a PTFE syringe filter (25 mm o.d., 0.45 μm pore disk filter; Chromatographic Specialties Inc.) and then combined. The combined extracts were then concentrated to ~ 0.1 mL under a gentle flow of analytical-grade nitrogen (Praxair Canada

isotope ratio mass spectrometer (IRMS; IsoPrime, GV Instruments, Manchester, U.K.) for isotope ratio and concentration measurements. The majority of extracts were analyzed on the day of preparation; otherwise, the extracts were stored in a refrigerator. All extracts were analyzed within 1 week of extraction. This methodology was thoroughly evaluated using 11 reference phenolic compounds,¹² and recovery yields of these compounds were better than 80%. Their retention times in the GCC–IRMS chromatograms and reference $\delta^{13}\text{C}$ values obtained using off-line combustion and dual-inlet isotope ratio analysis are listed in the Supporting Information, Table S-1.

The BSTFA derivatization (referred to above) replaces an active hydrogen atom of the OH group with a trimethylsilyl (TMS) group, causing a defined isotopic shift from the original $\delta^{13}\text{C}$ value. This isotopic shift was corrected as follows to obtain the $\delta^{13}\text{C}$ values of the TMS-free compounds ($\delta^{13}\text{C}_{\text{free}}$):

$$\delta^{13}\text{C}_{\text{free}} = \frac{\#C_{\text{deriv}}}{\#C_{\text{free}}} \delta^{13}\text{C}_{\text{deriv}} - \frac{\#C_{\text{TMS}}}{\#C_{\text{free}}} \delta^{13}\text{C}_{\text{TMS}} \quad (1)$$

where $\delta^{13}\text{C}_{\text{deriv}}$, $\delta^{13}\text{C}_{\text{TMS}}$, $\#C_{\text{deriv}}$, $\#C_{\text{TMS}}$, and $\#C_{\text{free}}$ are the $\delta^{13}\text{C}$ values for the derivatized compound, the $\delta^{13}\text{C}$ value for the TMS group, the total number of carbon atoms in the derivatized molecule, the number of carbon atoms in the TMS group, and the number of carbon atoms in the TMS-free compound. The carbon atom numbers were determined according to the stoichiometry of the derivatization reaction as long as the original compound was identified. We used $-49.94\% \pm 0.33\%$ (\pm standard error) for $\delta^{13}\text{C}_{\text{TMS}}$ because the BSTFA used here was the same batch as reported previously.¹³

3. RESULTS AND DISCUSSION

3.1. Product Identification. GCC–IRMS analysis of the PTFE blank filter extracts showed some contaminants in the extracts. Although the retention times of the contaminants differed from those of the target compounds, they differed from the retention times of the internal standards IS3 and IS4 by only ~ 0.4 min in some cases. The contaminant levels varied from sample to sample, and the measurement results for these internal standards may be influenced to some extent by these contaminant peaks.

GCC–IRMS analysis revealed significant abundances of eight product peaks in the flow reactor samples (Figure 1), while only one prominent product peak was observed in one of the smog chamber samples. We identified seven products based on GC–MS spectral patterns, retention times, and the abundance of the m/z 46 peak in the GCC–IRMS spectrum. The m/z 46 peak is an indication that the product contains a




Figure 1. Chromatogram of the m/z 44 trace obtained based on GCC–IRMS analysis of an SOA extract. IS1, IS2, IS3, IS4, C₁₇, C₁₈, and C₁₉ are catechol, 2-nitrophenol, 4-ethylresorcinol, 2,6-dimethyl-4-nitrophenol, heptadecane, octadecane, and nonadecane, used as internal or volumetric standards. The numbered peaks are products found in the SOA extracts. See Table 2 for product identification.

Table 2. Identified Products and Methods of Their Identification

peak no.	retention time (min)	identified product (abbrev)	method of identification ^a
1	48.9	4-nitrophenol (4NP)	MS and m/z 46
2	51.4	3-methyl-4-nitrophenol (3Me4)	RT, MS, and m/z 46
3	54.0	2-methyl-4-nitrophenol (2Me4)	RT, MS, and m/z 46
4	60.7	methylnitrocatechol isomer 1 (MeNC1)	MS and m/z 46
5	62.5	unknown	unsuccessful
6	63.0	methylnitrocatechol isomer 2 (MeNC2)	MS and m/z 46
7	63.4	methylnitrocatechol isomer 3 (MeNC3)	MS and m/z 46
8	64.8	nitrocatechol isomer (NC)	MS and m/z 46

^aRT, MS, and m/z 46 denote the identification method of retention time matching, mass spectral pattern, and detection of the high signal peak in the m/z 46 trace chromatogram (identification of the nitro group) based on GCC–IRMS analysis, respectively.

ally, a high abundance in the m/z 46 trace in the GCC–IRMS spectrum was observed for peak 1, suggesting that the product contained at least one nitro group. Based on this, peak 1 was most likely a structural isomer of nitrophenol. Analysis of 3- and 2-nitrophenols revealed that their retention times did not match the retention time for peak 1. Therefore, we identified peak 1 as 4-nitrophenol.

(*MM* = 313) and peak 8 as a nitrocatechol isomer (*MM* = 298). Unfortunately, peak 5 could not be identified.

3.2. Yields and $\delta^{13}\text{C}$ Values of Specific Products in SOA. $\delta^{13}\text{C}_{\text{free}}$ values and yields of identified products in the flow reactor samples for various extents of the toluene reaction are listed in Table 3. All $\delta^{13}\text{C}_{\text{free}}$ values were calculated according to eq 1 with the reference $\delta^{13}\text{C}_{\text{TMS}}$ value \pm standard error of $-49.94\text{‰} \pm 0.33\text{‰}$ determined for the TMS group of BSTFA in this batch.¹⁹ Yield is defined as the carbon mass of the identified product relative to the carbon mass of toluene reacted. Carbon masses were determined according to the internal calibration for structurally similar reference compounds listed in the Supporting Information, Table S-1. The yield measurements of the flow reactor samples showed that 2-methyl-4-nitrophenol accounted for 0.3–3% of reacted toluene, followed by methylnitrocatechol isomer 3 (0.1–0.6%), 4-nitrophenol and methylnitrocatechol isomer 1 (0.1–0.2%), and other products (<0.1%). These yields increased as the extent of toluene reaction progressed. Based on the dependence between total SOA carbon yields and the extent of toluene reaction processing,¹⁰ carbon mass fractions of 4-nitrophenol, 3-methyl-4-nitrophenol, 2-methyl-4-nitrophenol, methylnitrocatechol isomer 1, methylnitrocatechol isomer 2, methylnitrocatechol isomer 3, and nitrocatechol in SOA were $3.1\% \pm 0.5\%$, 3.2% (no error available), $54\% \pm 19\%$, $1.2\% \pm 0.2\%$, 0.9% (no error available), $7.6\% \pm 1\%$, and $0.5\% \pm 0.1\%$, respectively. The $\delta^{13}\text{C}_{\text{free}}$ values for all products, with the exception of 4-nitrophenol, in the flow reactor samples were between -29 and -37‰ , indicating that these products were significantly depleted in ^{13}C relative to the starting material, toluene ($\delta^{13}\text{C}$ for toluene $\approx -27\text{‰}$). In contrast, the $\delta^{13}\text{C}_{\text{free}}$ values for 4-nitrophenol ranged from -25 to -28‰ , indicating that ^{13}C in 4-nitrophenol was slightly enriched or similar to that in initial toluene. Comparison of photolysis rates of nitrophenol (3×10^{-4} – $7 \times 10^{-4} \text{ s}^{-1}$)²² with the product of the rate constants for photooxidation of nitrophenols ($\sim 5 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)²² multiplied by the estimated OH concentration in the flow reactor based on the toluene loss (1×10^9 – $1 \times 10^{10} \text{ molecules cm}^{-3}$) and the residence time (0.6–3.2 min)¹⁰ suggests that the photolysis accounted for less than 10%. Therefore, the wide range of observed $\delta^{13}\text{C}$ suggests that the products and their precursors underwent chemical reactions with different carbon KIEs.

Due to the very low nitrophenol concentrations, the analysis of smog chamber samples resulted in only one successful detection of 2-methyl-4-nitrophenol from the SCII experiment. Determined yield and $\delta^{13}\text{C}_{\text{free}}$ values were 0.042% and -29.3‰ , respectively. Compared to the total SOA carbon yields for the smog chamber experiments, 2-methyl-4-nitrophenol accounted for 0.042% of SOA carbon. This is

Table 3. Yields (\pm Standard Deviation from Triplicate Measurements) of Specific Products Identified in the SOA from Flow Reactor Experiments

Product	FR2 27 \pm 2% toluene reaction		FR3 21 \pm 2% toluene reaction		FR4 10 \pm 1% toluene reaction		FR5 9 \pm 1% toluene reaction	
	$\delta^{13}\text{C}$ (‰)	carbon yield (%)	$\delta^{13}\text{C}$ (‰)	carbon yield (%)	$\delta^{13}\text{C}$ (‰)	carbon yield (%)	$\delta^{13}\text{C}$ (‰)	carbon yield (%)
2-methyl-4-nitrophenol	-28 ± 1	0.170 ± 0.004	-25 ± 2	0.133 ± 0.008	-26.4 ± 0.3	0.019 ± 0.001	N/A ^b	N/A ^b
4-nitrophenol	-34.0 ± 0.2	0.083 ± 0.002	-32 ± 3	0.077 ± 0.006	N/A ^b	N/A ^b	N/A ^b	N/A ^b
3-methyl-4-nitrophenol	-30.5 ± 0.3	2.69 ± 0.02	N/A ^b	3.20 ^c	-33.1 ± 0.7	0.30 ± 0.01	-29.8 ± 0.4	0.256 ± 0.005
2-methyl-4-nitrophenol	-31.6 ± 0.6	0.110 ± 0.003	-30.9 ± 0.6	0.065 ± 0.002	-32.85 ± 0.01	0.015 ± 0.001	-32 ± 1	0.015 ± 0.001
methylnitrocatechol isomer 1	-32.4 ± 0.6	0.058 ± 0.002	-35.4 ± 0.9	0.041 ± 0.001	N/A ^b	N/A ^b	N/A ^b	N/A ^b
methylnitrocatechol isomer 2	-33.98^c	0.48 ± 0.01	-32.9 ± 0.3	0.28 ± 0.01	-32.7 ± 0.5	0.061 ± 0.001	-33.9 ± 0.3	0.064 ± 0.001
methylnitrocatechol isomer 3	-30.7 ± 0.6	0.051 ± 0.001	-32.1 ± 0.6	0.034 ± 0.001	-33.1 ± 1	0.022 ± 0.001	-32.4 ± 0.6	0.024 ± 0.001

^aof the identified product divided by the reacted carbon mass of toluene. ^bNo data available. ^cNo standard deviation is available due to only one available result.

Table 4. Difference^a (\pm Standard Error) between the Measured $\delta^{13}\text{C}_{\text{free}}$ Value for a Specific Product and Predicted $\delta^{13}\text{C}$ Value for the Sum of All Products ($\Delta \delta^{13}\text{C}$, ‰)

	FR1 0.17 fraction of toluene reacted	FR2 0.27 fraction of toluene reacted	FR3 0.21 fraction of toluene reacted	FR4 0.1 fraction of toluene reacted	FR5 0.09 fraction of toluene reacted
4NP	5.27 ± 0.49	3.84 ± 0.75	6.75 ± 0.92	5.75 ± 0.36	N/A ^b
3Me4	-4.53 ± 1.10	-2.36 ± 0.32	-0.36 ± 1.60	N/A ^b	N/A ^b
2Me4	-0.62 ± 0.33	1.33 ± 0.34	N/A ^b	-0.89 ± 0.50	2.40 ± 0.37
MeNC1	-0.20 ± 0.34	0.02 ± 0.44	0.93 ± 0.44	-0.67 ± 0.30	-0.22 ± 0.63
MeNC2	-1.77 ± 0.48	-0.78 ± 0.44	-3.53 ± 0.58	N/A ^b	N/A ^b
MeNC3	-0.57 ± 0.42	-2.35	-1.05 ± 0.34	-0.55 ± 0.42	-1.72 ± 0.34
NC	1.56 ± 0.66	0.91 ± 0.46	-0.29 ± 0.47	-0.93 ± 0.73	-0.23 ± 0.46

^a $\Delta \delta^{13}\text{C} = \delta^{13}\text{C}_{\text{free}} - \delta^{13}\text{C}$ for the sum of all products. ^bNot available.




Figure 2. Postulated reaction mechanisms for the photooxidation of toluene and the following oxidation reactions.^{16–19}

magnitude with standard errors based on triplicate measurements. The average $\Delta \delta^{13}\text{C}$ for each compound (excluding 4-nitrophenol and methylnitrocatechol isomer 2) over the five flow reactor experiments was not significantly different from zero at the 95% confidence level. Thus, we concluded that, for all products identified here (excluding 4-nitrophenol and methylnitrocatechol isomer 2), there was no significant difference between the $\delta^{13}\text{C}$ value predicted by a mass balance and the observed $\delta^{13}\text{C}$ value. In contrast, the $\Delta \delta^{13}\text{C}$ values for 4-nitrophenol were +6‰ with a standard deviation of 1‰, clearly showing a significant difference.

3.3. Comparison with Isotope Ratios Derived from Postulated Reaction Mechanisms. The majority of identified phenolic products have carbon isotope ratios that

proceeds through two channels: addition of OH to the aromatic ring (Figure 2, 1a channel), accounting for approximately 88% of the initial reaction, and H abstraction from the methyl group (Figure 2, 1b channel), accounting for approximately 12% of the initial reaction.²³ The sum of the second-order rate constants for the 1a and 1b channels is $5.96 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.²⁴ The 1a channel is known to form a methylhydroxycyclohexadienyl radical, which rapidly decomposes back to the reactants (lifetime of 0.3 s at 298 K and atmospheric pressure), but also undergoes several forward reactions (Figure 2, 2a, 2b, and 2c channels). The 2a channel is believed to yield benzoquinone and ring-opening products.²³ The 2b channel yields cresols (the first-generation product) with yields of 12–40% of reacted toluene.^{17,25–28} Furthermore, 1

channel is $\sim 4 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.³⁰ Under our experimental condition with an O_2 content of 20%, the 2c channel accounted for 30 and 80% of the loss of methylhydroxycyclohexadienyl radical at NO_2 mixing ratios of 1 ppmv (the condition in the smog chamber experiments) and 10 ppmv (the condition in the flow reactor experiments), respectively. The methylhydroxynitrocyclohexadienyl radical is believed to decompose and produce cresol and HONO .²⁷ Compared to OH addition, H abstraction from the OH substituent of cresol (3b channel) and the methyl substituent is a minor path ($\sim 7\%$) owing to the lower probability of OH attack on the OH substituent of cresol than OH attack on the aromatic ring, but still more favored than abstraction of an H atom from the methyl group.¹³ This was experimentally confirmed based on observations by Olariu et al.³¹ where hydroxybenzaldehyde, an expected product via H abstraction from the methyl substituent, was not formed. They proposed that the branching ratios of OH addition and H abstraction are 90 and 10%, respectively. The 3a and 3b channels yield methylcatechols and methylnitrophenols, respectively,³¹ which may further react with the OH radical. Rate constants for reactions of methylnitrophenols with the OH radical in the gas phase (4a and 4b channels in Figure 3) range from 3×10^{-12} to

experimentally determined overall KIE for the reaction of toluene with OH radical (${}^{\text{OH}}\epsilon_{\text{C}_7\text{H}_8}$) was $5.95\% \pm 0.28\%$.³³

Note that all values of KIE stated herein are expressed as epsilon values ($\epsilon = k_{12}/k_{13} - 1$, where k_{12} is the rate constant for a reactant containing only ^{12}C carbons in the molecule and k_{13} is the rate constant for a reactant containing both ^{12}C carbons and a ^{13}C carbon in the molecule). The ${}^{\text{OH}}\epsilon_{\text{C}_7\text{H}_8}$ value is the sum of the two channel-specific KIEs:

$${}^{\text{OH}}\epsilon_{\text{C}_7\text{H}_8} = {}^{\text{OH}}R_{\text{adct}} \frac{1}{{}^{\text{OH}}N_{\text{C}}} {}^{\text{OH}}\epsilon_{\text{adct}} + {}^{\text{OH}}R_{\text{abst}} \frac{1}{{}^{\text{OH}}N_{\text{C}}} {}^{\text{OH}}\epsilon_{\text{abst}} \quad (2)$$

where ${}^{\text{OH}}R_{\text{abst}}$, ${}^{\text{OH}}\epsilon_{\text{abst}}$, ${}^{\text{OH}}R_{\text{adct}}$, ${}^{\text{OH}}\epsilon_{\text{adct}}$ and N_{C} are the branching ratio for H abstraction from a methyl group, the primary carbon KIE for H abstraction from a ^{13}C atom in the methyl group, the branching ratio for OH addition to the aromatic carbon ring, the primary carbon KIE for OH addition to a ^{13}C atom in the aromatic ring, and the carbon number of reactant molecules, respectively. Note that the KIEs over the N_{C} times the branching ratios indicate channel-specific primary carbon KIEs, in which the probability of reaction at a ^{13}C atom was considered. Methods to estimate the channel-specific KIEs assume no site preference in the distribution of the ^{13}C atom within the molecule, which has been validated according to the results of structure-related reactivity analysis as well as systematically changing carbon KIEs (i.e., agreement with the trend of $1/N_{\text{C}}$) for groups of *n*-alkanes or aromatic hydrocarbons.^{33,34} Using eq 2 with the ${}^{\text{OH}}\epsilon_{\text{abst}}$ of $18.7\% \pm 5.2\%$ ³⁴ and the aforementioned ${}^{\text{OH}}\epsilon_{\text{C}_7\text{H}_8}$, the ${}^{\text{OH}}\epsilon_{\text{adct}}$ for OH addition to the toluene ring was calculated to be $44.2\% \pm 5.2\%$. This value agrees with the uncertainty range of the estimated ${}^{\text{OH}}\epsilon_{\text{adct}}$ of $40.6\% \pm 1.4\%$ from the best-fit curve of KIEs versus the number of carbon atoms for a variety of aromatic hydrocarbons.³³ ${}^{\text{OH}}\epsilon_{\text{adct}}$ or ${}^{\text{OH}}\epsilon_{\text{abst}}$ normalized by probabilities of ^{13}C distributions in the methyl group and the aromatic ring allows a probability-based KIE to be calculated for each channel of the toluene reaction with OH. By multiplying these values by the branching ratios and the channel-specific k_{12} values, channel-specific k_{13} values were obtained. In addition to the ${}^{\text{OH}}\epsilon_{\text{abst}}$ and the ${}^{\text{OH}}\epsilon_{\text{adct}}$ there was a channel-specific KIE for H abstraction from the OH groups of hydroxyaromatic compounds. We assumed this KIE to be negligibly small ($\sim 0\%$) since its impact on $\delta^{13}\text{C}$ is a secondary isotope effect. Nevertheless, it cannot be excluded completely that the carbon KIE for this reaction is somewhat larger than expected for a typical secondary isotope effect due to delocalization of the unpaired electron of the phenoxy group. However, carbon KIEs for reactions of phenolic compounds with OH radicals have not been measured, and any assumption of a significant KIE would be




Figure 3. Production reaction mechanisms used to calculate $\delta^{13}\text{C}$ of methylnitrocatechol.

Table 5. Rate Constants, Kinetic Isotope Effects, and Branching Ratios Used for the Isotope Model

reaction channel	overall k ($\text{cm}^3 \text{s}^{-1} \text{molecule}^{-1}$)	branching ratio	channel-specific k_{12} ($\text{cm}^3 \text{s}^{-1} \text{molecule}^{-1}$)	channel-specific ϵ (%)	channel-specific k_{13} ($\text{cm}^3 \text{s}^{-1} \text{molecule}^{-1}$)
1a	5.96×10^{-12} ^a	0.88 ^a	5.2×10^{-12}	7.5 ^b	5.2×10^{-12}
1b		0.12 ^a	7.2×10^{-13}	0.27 ^b	7.1×10^{-13}
2a	$(5.4 \pm 0.6) \times 10^{-16}$ ^c	0.1	5×10^{-17}	0	5×10^{-16}
2b		0.9	4.9×10^{-16}	0	4.9×10^{-16}
2c	4.0×10^{-11} ^d	1.0	3.6×10^{-11}	0	3.6×10^{-11}
3a	4.10×10^{-11} ^d	0.9	4×10^{-11}	5.7 ^e	3.7×10^{-12}
3b		0.1	4×10^{-12}	0	4×10^{-12}
4a	5×10^{-12} ^f	0.9	5×10^{-12}	5.7 ^e	4.5×10^{-12}
4b		0.1	5×10^{-13}	0	5×10^{-13}
5a	2.05×10^{-11} ^g	0.9	1.9×10^{-11}	5.7 ^e	1.8×10^{-11}
5b		0.1	2×10^{-12}	0	2×10^{-12}

^aThe value was adapted from Atkinson.²⁴ ^bThe KIE reported by Anderson et al.³⁴ was used to estimate the channel-specific ϵ . ^cThe second-order k for the reaction of methylcyclohexadienyl radical with O_2 was adopted from Knispel et al.³⁰ ^dThe value is for the *o*-cresol + OH reaction in Calvert et al.²³ ^eThe KIE for this channel was estimated according to the generalized carbon KIE for OH adduct reaction to the aromatic ring ($40.6\% \pm 1.4\%$) divided by N_o , reported by Anderson et al.³³ ^fThe value was adapted from Bejan et al.²² ^gThe value was adapted from Olariu et al.³²

Predicted $\delta^{13}\text{C}$ values were calculated for methylnitrophenol, the second-generation product via the 3b channel, methylcyclohexadienyl radical, and the sum of all products from the photooxidation of toluene (Figure 4). The calculations were





Figure 4. Measured and predicted $\delta^{13}\text{C}$ values for methylnitrophenol (MeNP). As a comparison, predicted $\delta^{13}\text{C}$ values for the methylcyclohexadienyl radical and the sum of all products from toluene photooxidation are also shown.

made for two cases: case I, most of the produced methylnitrophenol partitioned into the condensed phase immediately and no further reaction takes place; case II, 90% of produced methylnitrophenol remained in the gas phase and undergoes further reactions, and the remaining methylnitrophenol partitions to the condensed phase, which is under equilibrium with methylnitrophenol in the gas phase. For case II, note that “under equilibrium” implies that $\delta^{13}\text{C}$ values of

with the predicted $\delta^{13}\text{C}$ values for methylnitrophenols. It remains unclear whether this discrepancy can be attributed to our assumptions on unavailable KIEs (i.e., 0% KIEs, as shown in Table 5) or whether it indicated likewise the results of yields that the production is via different reaction channels, such as the 2c channel proposed by Zellner et al.²⁹ It is possible that the NO_2 addition to methylcyclohexadienyl radical forms a NO_2 adduct with a high excitation state, which may undergo further reactions with OH, resulting in the formation of methylnitrophenols and methylnitrocatechols. At this time, experimental evidence to support this hypothesis is not available.

Assuming that the majority of methylnitrocatechol partitions into the condensed phase with no loss reaction in the condensed phase, $\delta^{13}\text{C}$ values for methylnitrocatechol were calculated for three cases: case A, it is produced via the 4a and the 5b channels; case B, it is produced via the 4a channel only; case C, it is produced via the 5b channel only (Figure 5). Figure 5 also shows predicted $\delta^{13}\text{C}$ values for the methylcyclohexadienyl radical, methylcatechol, and methylnitrophenol for comparison. Note that, in these calculations, the majority (90%) of the methylnitrocatechol precursor, methylnitrophenol, was assumed to react in the gas phase and produce particulate methylnitrocatechol. As discussed earlier, the



observed $\delta^{13}\text{C}_{\text{free}}$ values for methylnitrocatechol isomer 1 agree with the predicted $\delta^{13}\text{C}$ values for the sum of all products within 2σ , but disagree with the predicted $\delta^{13}\text{C}$ for methylnitrocatechol in all three cases. The observed $\delta^{13}\text{C}_{\text{free}}$ values for methylnitrocatechol isomers 2 and 3 show larger fluctuations, but are overall slightly lower than that for isomer 1 and closer to the predicted $\delta^{13}\text{C}$ for methylnitrocatechol. Unfortunately, the number of observations in this limited range of toluene turnover is insufficient to conclude whether they follow the predicted $\delta^{13}\text{C}$ trends for methylnitrocatechol.

Figure 6 shows the mechanism for production of nitrophenol proposed by Atkinson.¹⁴ For the production of 4-nitrophenol,




Figure 6. Postulated reaction mechanisms for production of nitrophenol via 1b channel.¹⁴

the reaction mechanism involves a reaction at the methyl group of toluene; therefore, production of 4-nitrophenol must proceed via the 1b channel in Figure 2. Subsequent reactions are shown in detail in Figure 6. According to the mechanism, significant carbon isotope fractionations are expected at the initial step (i.e., H abstraction from the methyl group) when benzaldehyde reacts with OH radical (i.e., abstraction of a hydrogen atom from the aldehyde group) because both reactions are cleavages of primary C–H bonds (carbon KIEs). These fractionations result in ^{13}C depletion in the reaction products. However, this carbon isotope fractionation




Figure 7. Measured $\delta^{13}\text{C}$ value for 4-nitrophenol and predicted $\delta^{13}\text{C}$ values for toluene, benzaldehyde, nitrophenol, and the sum of all products.

values for 4-nitrophenol were similar to the initial $\delta^{13}\text{C}$ of toluene, while the $\delta^{13}\text{C}$ values of the other phenolic compounds were depleted by 5–6‰ compared to the initial $\delta^{13}\text{C}$ value of toluene. For 4-nitrophenol, the $\delta^{13}\text{C}$ value dependence on precursor conversion was consistent with the formation mechanism proposed in previous reports. However, the observed $\delta^{13}\text{C}$ values for the other phenolic compounds were significantly lower than $\delta^{13}\text{C}$ values estimated for these products according to the postulated mechanisms (i.e., production via the 3a and 3b channels). Although the observed $\delta^{13}\text{C}$ values varied, they were similar to the predicted $\delta^{13}\text{C}$ values for the sum of all products using the mass balance. This suggests that these particulate phenolic products were produced via different reaction channels. To constrain production channels, more experimental evidence, such as KIE studies for intermediates, is required. Specifically, the lack of measured KIEs for reactions of phenols with OH radicals creates uncertainty in predicted $\delta^{13}\text{C}$ values for phenolic compounds by the oxidation of toluene in air. Nevertheless, our results demonstrate the usefulness of compound-specific stable carbon isotope measurements to increase our understanding of chemical reaction mechanisms.

ASSOCIATED CONTENT

Supporting Information

Retention time index and reference $\delta^{13}\text{C}$ values for standard compounds (Table S-1); GCC–IRMS chromatogram for a blank extract (Figure S-1). This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

ACKNOWLEDGMENTS

We thank Darrell Ernst, Alina Chivulescu, and Wendy Zhang at Environment Canada for technical support. We also acknowledge the Natural Sciences and Engineering Research Council of Canada, the Canadian Foundation for Climate and Atmospheric Sciences, and Environment Canada for financial support. The project was also partially supported by the ESPEC Foundation for Global Environment Research and Technologies (Charitable Trust).

REFERENCES

- (1) Jimenez, J. L.; Canagaratna, M. R.; Donahue, N. M.; Prevot, A. S. H.; Zhang, Q.; Kroll, J. H.; DeCarlo, P. F.; Allan, J. D.; Coe, H.; Ng, N. L.; et al. Evolution of Organic Aerosols in the Atmosphere. *Science* **2009**, *326*, 1525–1529.
- (2) Dockery, D. W.; Pope, C. A., 3rd; Xu, X.; Spengler, J. D.; Ware, J. H.; Fay, M. E.; Ferris, B. G., Jr.; Speizer, F. E. An Association between Air Pollution and Mortality in Six U.S. Cities. *New Engl. J. Med.* **1993**, *329*, 1753–1759.
- (3) Anderson, T. L.; Charlson, R. J.; Schwartz, S. E.; Knutti, R.; Boucher, O.; Rodhe, H.; Heintzenberg, J. Atmospheric Science: Climate Forcing by Aerosols □ A Hazy Picture. *Science* **2003**, *300*, 1103–1104.
- (4) Rudolph, J. *Volatile Organic Compounds in the Atmosphere*; Blackwell Publishing: Oxford, U.K., 2007.
- (5) Sakugawa, H.; Kaplan, I. R. Stable Carbon Isotope Measurements of Atmospheric Organic Acids in Los Angeles, California. *Geophys. Res. Lett.* **1995**, *22*, 1509–1512.
- (6) Moukhtar, S.; Saccon, M.; Kornilova, A.; Irei, S.; Huang, L.; Rudolph, J. Method for Determination of Stable Carbon Isotope Ratio of Methylnitrophenols in Atmospheric PM. *Atmos. Meas. Technol.* **2011**, *4*, 2453–2464.
- (7) Wang, H.; Kawamura, K. Stable Carbon Isotopic Composition of Low-Molecular Weight Dicarboxylic Acids and Ketoacids in Remote Marine Aerosol. *J. Geophys. Res. Atmos.* **2006**, *111*, D07304.
- (8) Gensch, I.; Kiendler-Scharr, A.; Rudolph, J. Isotope Ratio Studies of Atmospheric Organic Compounds: Principles, Methods, Applications, and Potential. *Int. J. Mass Spectrom.* **2014**, *365–366*, 206–221.
- (9) Irei, S.; Takami, A.; Hayashi, M.; Sadanaga, Y.; Hara, K.; Kaneyasu, N.; Sato, K.; Arakaki, T.; Hatakeyama, S.; Bandow, H.; Hikida, T.; Shimono, A. Transboundary Secondary Organic Aerosol in Western Japan Indicated by the $\delta^{13}\text{C}$ of Water-Soluble Organic Carbon and the m/z 44 signal in Organic Aerosol Mass Spectra. *Environ. Sci. Technol.* **2014**, *48*, 6273–6281.
- (10) Irei, S.; Huang, L.; Collin, F.; Zhang, W.; Hastie, D.; Rudolph, J. Flow Reactor Studies of the Stable Carbon Isotope Composition of Secondary Particulate Organic Matter Generated by OH-Radical-Induced Reactions of Toluene. *Atmos. Environ.* **2006**, *40*, 5858–5867.
- (11) Irei, S.; Rudolph, J.; Huang, L.; Auld, J.; Hastie, D. Stable Carbon Isotope Ratio of Secondary Particulate Organic Matter Formed by Photooxidation of Toluene in Indoor Smog Chamber. Situ Determination of Ring-Retaining Product Yields. *J. Phys. Chem. A* **1998**, *102*, 10289–10299.
- (16) Forstner, H. J. L.; Flagan, R. C.; Seinfeld, J. H. Secondary Organic Aerosol from the Photooxidation of Aromatic Hydrocarbons: Molecular Composition. *Environ. Sci. Technol.* **1997**, *31*, 1345–1358.
- (17) Jang, M.; Kamens, R. M. Characterization of Secondary Aerosol from the Photooxidation of Toluene in the Presence of NO_x and 1-Propene. *Environ. Sci. Technol.* **2001**, *35*, 3626–3639.
- (18) Sato, K.; Hatakeyama, S.; Imamura, T. Secondary Organic Aerosol Formation during the Photooxidation of Toluene: NO_x Dependence of Chemical Composition. *J. Phys. Chem. A* **2007**, *111*, 9796–9808.
- (19) Bennett, J. F.; Collin, F.; Hastie, D. R. A Laboratory Flow Reactor with Gas Particle Separation with On-Line MS/MS for Product Identification in Atmospherically Important Reactions. *Atmos. Meas. Technol.* **2009**, *2*, 813–823.
- (20) Auld, J.; Hastie, D. R. Investigation of Organic Nitrate Product Formation during Hydroxy Radical Initiated Photo-Oxidation of β -Pinene. *Atmos. Environ.* **2011**, *45*, 26–34.
- (21) Irei, S. Ph.D. Dissertation, York University, Toronto, Canada, 2010.
- (22) Bejan, I.; Barnes, I.; Olariu, R.; Zhou, S.; Wiesen, P.; Benter, T. Investigations on the Gas-Phase Photolysis and OH Radical Kinetics of Methyl-2-nitrophenols. *Phys. Chem. Chem. Phys.* **2007**, *9*, 5686–5692.
- (23) Calvert, J. G.; Atkinson, R.; Becker, K. H.; Kamens, R. M.; Seinfeld, J. H.; Wallington, T. H.; Yarwood, G. *The Mechanisms of Atmospheric Oxidation of Aromatic Hydrocarbons*; Oxford University Press: New York, 2002.
- (24) Atkinson, R. Gas-Phase Tropospheric Chemistry of Organic Compounds. *J. Phys. Chem. Ref. Data, Monogr.* **1994**, No. No. 2, 1–216.
- (25) Gery, M. W.; Fox, D. L.; Jeffries, H. E.; Stockburger, L.; Weathers, W. S. A Continuous Stirred Tank Reactor Investigation of the Gas-Phase Reaction of Hydroxyl Radicals and Toluene. *Int. J. Chem. Kinet.* **1985**, *17*, 931–955.
- (26) Leone, J. A.; Seinfeld, J. H. Comparative Analysis of Chemical Reaction Mechanisms for Photochemical Smog. *Atmos. Environ., Part A: Gen. Top.* **1985**, *19*, 437–464.
- (27) Atkinson, R.; Aschmann, S. M. Products of the Gas-Phase Reactions of Aromatic Hydrocarbons: Effect of NO_2 Concentration. *Int. J. Chem. Kinet.* **1994**, *26*, 929–944.
- (28) Smith, D. F.; McIver, C. D.; Kleindienst, T. E. Primary Product Distribution from the Reaction of Hydroxyl Radicals with Toluene at ppb NO_x Mixing Ratios. *J. Atmos. Chem.* **1998**, *30*, 209–228.
- (29) Zellner, R.; Fritz, B.; Preidel, M. A cw UV Laser Absorption Study of the Reactions of the Hydroxy-Cyclohexadienyl Radical with NO_2 and NO . *Chem. Phys. Lett.* **1985**, *121*, 412–416.
- (30) Knispel, R.; Koch, R.; Siese, M.; Zetzsch, C. Adduct Formation of OH Radicals with Benzene, Toluene, and Phenol and Consecutive Reactions of the Adducts with NO_x and O_2 . *Ber. Bunsen-Ges. Phys. Chem.* **1990**, *94*, 1375–1379.
- (31) Olariu, R. I.; Klotz, B.; Barnes, I.; Becker, K. H.; Mocanu, R. FT-IR Study of the Ring-Retaining Products from the Reaction of OH Radicals with Phenol, *o*-, *m*-, and *p*-Cresol. *Atmos. Environ.* **2002**, *36*, 2695–2697.

